

# Multiple Indicator Cluster Survey

## Model Full Report

### TABLES

<a href="#">Table 1: Number of households and women, and response rates, Country, Year</a>	3
<a href="#">Table 2: Single year age distribution of household population by sex, Country, Year</a>	4
<a href="#">Table 3: Percentage of cases with missing information, Country, Year</a>	5
<a href="#">Table 4: Percent distribution of households by background characteristics, Country, Year</a>	6
<a href="#">Table 5: Percent distribution of women 15-49 by background characteristics, Country, Year</a>	7
<a href="#">Table 6: Percent distribution of children under 5 by background characteristics, Country, Year</a>	8
<a href="#">Table 7: Mean number of children ever born (CEB) and proportion dead by mother's age, Country, Year</a>	9
<a href="#">Table 8: infant and under-five mortality rates</a>	10
<a href="#">Table 9: Percentage of children aged 36-59 months who are attending some form of organized early childhood education programme, Country, Year</a>	11
<a href="#">Table 10: Percentage of children of primary school age attending primary school, Country, Year</a>	12
<a href="#">Table 11: Percentage of children entering first grade of primary school who eventually reach grade 5, Country, Year</a>	13
<a href="#">Table 12: Percentage of the population aged 15 years and older that is literate, Country, Year</a>	14
<a href="#">Table 13: Percentage of the population using improved drinking water sources, Country, Year</a>	15
<a href="#">Table 14: Percentage of the population using sanitary means of excreta disposal, Country, Year</a>	16
<a href="#">Table 15: Percentage of under-five children who are severely or moderately undernourished, Country, Year</a>	17
<a href="#">Table 16: Percent of living children by breastfeeding status, Country, Year</a>	18
<a href="#">Table 17: Percentage of households consuming adequately iodized salt, Country, Year</a>	19
<a href="#">Table 18: Percent distribution of children aged 6-59 months by whether they have received a high dose Vitamin A supplement in the last 6 months, Country, Year</a>	20
<a href="#">Table 19: Percentage of women with a birth in the last 12 months by whether they received a high dose Vitamin A supplement before the infant was 8 weeks old, Country, Year</a>	21
<a href="#">Table 20: Percentage of live births in the last 12 months that weighed below 2500 grams at birth, Country, Year</a>	22

<a href="#">Table 21: Percentage of children age 12-23 months immunized against childhood diseases at any time before the survey and before the first birthday, Country, Year</a> .....	23
<a href="#">Table 22: Percentage of children age 12-23 months currently vaccinated against childhood diseases, Country, Year</a> .....	24
<a href="#">Table 23: Percentage of under-five children with diarrhea in the last two weeks and treatment with ORS or ORT, Country, Year</a> .....	25
<a href="#">Table 24: Percentage of under-five children with diarrhea in the last two weeks who took increased fluids and continued to feed during the episode, Country, Year</a> .....	26
<a href="#">Table 25: Percentage of under-five children with acute respiratory infection in the last two weeks and treatment by health providers, Country, Year</a> .....	27
<a href="#">Table 26: Percentage of children 0-59 months of age reported ill during the last two weeks who received increased fluids and continued feeding, Country, Year</a> .....	28
<a href="#">Table 27: Percentage of caretakers of children 0-59 months who know at least 2 signs for seeking care immediately, Country, Year</a> .....	29
<a href="#">Table 28: Percentage of children 0-59 months of age who slept under an insecticide-impregnated bednet during the previous night, Country, Year</a> .....	30
<a href="#">Table 29: Percentage of children 0-59 months of age who were ill with fever in the last two weeks who received anti-malarial drugs, Country, Year</a> .....	31
<a href="#">Table 30: Percentage of women aged 15-49 who know the main ways of preventing HIV transmission, Country, Year</a> .....	32
<a href="#">Table 31: Percentage of women aged 15-49 who correctly identify misconceptions about HIV/AIDS, Country, Year</a> .....	33
<a href="#">Table 32: Percentage of women aged 15-49 who correctly identify means of HIV transmission from mother to child, Country, Year</a> .....	34
<a href="#">Table 33: Percentage of women aged 15-49 who express a discriminatory attitude towards people with HIV/AIDS, Country, Year</a> .....	35
<a href="#">Table 34: Percentage of women aged 15-49 who have sufficient knowledge of HIV/AIDS transmission, Country, Year</a> .....	36
<a href="#">Table 35: Percentage of women aged 15-49 who know where to get an AIDS test and who have been tested, Country, Year</a> .....	37
<a href="#">Table 36: Percentage of married or in union women aged 15-49 who are using (or whose partner is using) a contraceptive method, Country, Year</a> .....	38
<a href="#">Table 37: Percentage of mothers with a birth in the last 12 months protected against neonatal tetanus, Country, Year</a> .....	39
<a href="#">Table 38: Percent distribution of women aged 15-49 with a birth in the last year by type of personnel delivering antenatal care, Country, Year</a> .....	40
<a href="#">Table 39: Percent distribution of women aged 15-49 with a birth in the last year by type of personnel assisting at delivery, Country, Year</a> .....	41
<a href="#">Table 40: Percent distribution of children aged 0-59 months by whether birth is registered and reasons for non-registration, Country, Year</a> .....	42
<a href="#">Table 41: Percentage of children 0-14 years of age in households not living with a biological parent, Country, Year</a> .....	43
<a href="#">Table 42: Percentage of children 5-14 years of age who are currently working, Country, Year</a> .....	44

**Table 1: Number of households and women, and response rates, Country, Year**

	<b>Urban</b>	<b>Rural</b>	<b>Total</b>
Sampled households	2595	2076	4671
Occupied households	2535	2042	4577
Interviewed households	2307	1978	4285
Household response rate	91.0	96.9	93.6
Eligible women	2328	2080	4236
Interviewed women	2060	1905	3965
Women response rate	88.5	91.6	90.0
Children under 5	926	1031	1957
Interviewed children under 5	857	974	1831
Child response rate	92.6	94.5	93.6

**Table 2: Single year age distribution of household population by sex, Country, Year**

Age	Male		Female		Age	Male		Female	
	Number	Percent	Number	Percent		Number	Percent	Number	Percent
0	214	2.5	188	2.2	37	104	1.2	114	1.3
1	196	2.3	187	2.2	38	125	1.5	120	1.4
2	259	3.1	188	2.2	39	133	1.6	113	1.3
3	176	2.1	191	2.3	40	126	1.5	124	1.5
4	174	2.0	163	1.9	41	90	1.1	110	1.3
5	154	1.8	157	1.9	42	110	1.3	79	.9
6	176	2.1	158	1.9	43	82	1.0	99	1.2
7	193	2.3	160	1.9	44	88	1.0	84	1.0
8	173	2.0	147	1.7	45	106	1.3	101	1.2
9	180	2.1	175	2.1	46	56	.7	68	.8
10	195	2.3	172	2.0	47	76	.9	88	1.0
11	156	1.8	172	2.0	48	66	.8	89	1.1
12	177	2.1	150	1.8	49	55	.7	65	.8
13	177	2.1	201	2.4	50	76	.9	82	1.0
14	183	2.2	193	2.3	51	41	.5	52	.6
15	192	2.3	221	2.6	52	63	.7	46	.5
16	164	1.9	204	2.4	53	51	.6	48	.6
17	184	2.2	192	2.3	54	51	.6	70	.8
18	177	2.1	140	1.7	55	55	.6	63	.7
19	155	1.8	163	1.9	56	40	.5	53	.6
20	169	2.0	175	2.1	57	39	.5	49	.6
21	141	1.7	134	1.6	58	46	.5	53	.6
22	142	1.7	146	1.7	59	34	.4	56	.7
23	140	1.6	128	1.5	60	58	.7	61	.7
24	116	1.4	131	1.6	61	44	.5	46	.5
25	140	1.7	152	1.8	62	34	.4	44	.5
26	124	1.5	116	1.4	63	45	.5	49	.6
27	120	1.4	132	1.6	64	39	.5	40	.5
28	145	1.7	129	1.5	65	53	.6	61	.7
29	119	1.4	134	1.6	66	31	.4	42	.5
30	171	2.0	159	1.9	67	46	.5	44	.5
31	116	1.4	113	1.3	68	33	.4	34	.4
32	143	1.7	155	1.8	69	34	.4	35	.4
33	126	1.5	134	1.6	70+	261	3.1	294	3.5
34	120	1.4	116	1.4	Missing/DK	136	1.6	50	.6
35	158	1.9	139	1.6					
36	128	1.5	119	1.4	Total	8505	100.0	8457	100.0

**Table 3: Percentage of cases with missing information, Country, Year**

	<b>Reference population</b>	<b>Percent missing</b>	<b>Number</b>
Level of education	Household members	0.6	13724
Year of education	Household members	3.0	13723
Number of hours worked	Working children age 5-14	3.3	59
Complete birth date	Women 15-49	0.2	3965
Date of last tetanus toxoid injection	Women with a live birth in the last year	3.3	387
Ever been tested for HIV	Women 15-49	0.9	3707
Complete birth date	Children under 5	0.4	1830
Diarrhoea in last 2 weeks	Children under 5	1.6	1830
Weight	Children under 5	5.1	1830
Height	Children under 5	5.1	1830

**Table 4: Percent distribution of households by background characteristics, Country, Year**

	Area		Total
	Urban	Rural	
Central	74.3	0.0	41.5
South Central	17.1	27.9	21.9
West	8.6	24.8	15.7
East	0.0	26.1	11.5
South	.0.0	21.2	9.4
Number of HH members			
1	9.8	10.5	10.1
2-3	35.6	33.0	34.4
4-5	36.5	34.2	35.5
6-7	12.8	15.1	13.8
8-9	4.0	5.2	4.5
10+	1.4	2.1	1.7
Total	100.0	100.0	100.0
At least one child age < 15	60.9	66.1	63.2
At least one child age < 5	34.7	39.8	37.0
At least one woman age 15-49	78.9	79.5	79.2
Number	2395	1890	4285
Unweighted	2307	1978	4285

**Table 5: Percent distribution of women 15-49 by background characteristics, Country, Year**

	Area		Total
	Urban	Rural	
Central	72.2	0.0	40.0
South Central	19.0	30.3	24.0
West	8.8	24.6	15.8
East	0.0	24.7	11.0
South	0.0	20.4	9.1
Age			
15-19	18.2	23.6	20.6
20-24	16.4	15.3	15.9
25-29	15.0	14.9	15.0
30-34	15.4	14.7	15.1
35-39	13.9	12.8	13.4
40-44	11.9	10.1	11.1
45-49	9.1	8.5	8.8
Marital status			
Currently married	62.3	66.5	64.2
Not currently married	37.7	33.5	35.8
Ever given birth			
Yes	66.2	67.9	67.0
No	33.8	32.1	33.0
Education level			
None	2.3	12.2	6.7
Primary	19.2	35.5	26.5
Secondary +	78.5	52.3	66.8
Total	100.0	100.0	100.0
Number	2196	1769	3965
Unweighted	2060	1905	3965

**Table 6: Percent distribution of children under 5 by background characteristics, Country, Year**

	Area		Total
	Urban	Rural	
Male	52.2	53.8	53.0
Female	47.8	46.2	47.0
Central	74.0	.0	36.2
South Central	17.4	22.9	20.2
West	8.6	20.6	14.8
East	.0	23.4	11.9
South	.0	33.1	16.9
Age			
< 6 months	12.2	11.4	11.8
6-11 months	8.1	9.1	8.6
12-23 months	19.1	21.0	20.0
24-35 months	25.1	21.2	23.1
36-47 months	18.7	19.0	18.9
48-59 months	16.8	18.3	17.6
Mother's education			
None	5.5	20.2	13.0
Primary	16.9	36.9	27.1
Secondary +	77.6	42.9	59.9
Total	100.0	100.0	100.0
Number	896	934	1830
Unweighted	857	974	1831

**Table 7: Mean number of children ever born (CEB) and proportion dead by mother's age, Country, Year**

	<b>Mean number of CEB</b>	<b>Proportion dead</b>	<b>Number of women</b>
15-19	.162	.058	818
20-24	.925	.059	631
25-29	1.737	.032	593
30-34	2.547	.053	599
35-39	3.192	.047	531
40-44	3.718	.054	441
45-49	4.179	.079	351
Total	2.037	.055	3965

**Table 8: infant and under-five mortality rates**

---

	<b>Infant mortality rate</b>	<b>Under-five mortality rate</b>
Total	45 per 1000	52 per 1000

---

Reference date is 1992.9

**Table 9: Percentage of children aged 36-59 months who are attending some form of organized early childhood education programme, Country, Year**

	Attending programme	Number of children
Male	40.9	339
Female	39.2	327
Central	54.6	240
South Central	43.0	133
West	45.8	116
East	31.0	81
South	1.0	97
Urban	52.3	318
Rural	28.9	348
36-47 months	16.1	345
48-59 months	65.8	321
Mother's education		
None	16.2	80
Primary	30.3	197
Secondary +	49.9	389
<b>Total</b>	<b>40.1</b>	<b>666</b>

**Table 10: Percentage of children of primary school age attending primary school, Country, Year**

	Male		Female		Total	
	Attending	Number	Attending	Number	Attending	Number
Central	96.5	504	97.3	456	96.9	959
South Central	92.5	324	96.1	296	94.2	620
West	89.7	201	91.1	211	90.4	412
East	93.0	203	87.0	169	90.3	373
South	50.6	172	53.6	159	52.0	331
Urban	95.4	702	96.7	627	96.0	1329
Rural	81.5	702	82.3	663	81.9	1365
Age						
5	83.7	154	81.9	157	82.8	311
6	83.8	176	86.4	158	85.1	334
7	87.8	193	89.2	160	88.4	353
8	91.1	173	92.8	147	91.9	320
9	88.7	180	91.8	175	90.2	355
10	92.0	195	91.6	172	91.8	366
11	90.1	156	90.5	172	90.3	328
12	89.7	177	89.6	150	89.7	327
<b>Total</b>	<b>88.5</b>	<b>1404</b>	<b>89.3</b>	<b>1290</b>	<b>88.9</b>	<b>2694</b>

**Table 11: Percentage of children entering first grade of primary school who eventually reach grade 5, Country, Year**

	<b>Percent in grade 1 eventually reaching grade 2</b>	<b>Percent in grade 2 eventually reaching grade 3</b>	<b>Percent in grade 3 eventually reaching grade 4</b>	<b>Percent in grade 4 eventually reaching grade 5</b>	<b>Percent who reach grade 5 of those who enter grade 1</b>
Male	86.0	92.4	93.2	93.2	69.1
Female	86.1	92.8	95.2	89.5	68.1
Central	98.2	.	96.9	96.3	.
South Central	93.5	94.5	93.9	96.1	79.7
West	82.5	90.1	93.6	83.4	58.0
East	86.6	92.9	95.4	89.6	68.7
South	51.8	76.4	80.6	72.4	23.1
Urban	95.4	98.7	96.8	96.0	87.4
Rural	78.8	87.4	91.6	86.1	54.3
Total	86.1	92.6	94.2	91.4	68.6

**Table 12: Percentage of the population aged 15 years and older that is literate, Country, Year**

	Male			Female			Total		
	Literate	Not known	Number	Literate	Not known	Number	Literate	Not known	Number
Central	95.2	.6	2394	93.1	.6	2409	94.2	.6	4803
South Central	91.9	.9	1357	88.7	.2	1311	90.3	.6	2668
West	85.5	5.5	979	83.9	4.8	912	84.7	5.2	1892
East	91.1	.8	679	82.5	.7	666	86.9	.8	1345
South	54.1	13.7	175	38.2	7.9	507	42.3	9.4	682
Urban	94.1	1.0	3310	91.8	.8	3267	92.9	.9	6577
Rural	86.4	3.4	2274	75.5	3.1	2539	80.7	3.2	4814
Age									
15-24	93.2	2.7	1580	93.3	.8	1634	93.2	1.7	3214
25-34	94.9	1.4	1325	91.7	1.3	1338	93.3	1.4	2663
35-44	93.9	1.3	1145	88.2	1.5	1100	91.1	1.4	2245
45-54	90.2	1.2	642	78.5	2.5	709	84.0	1.9	1350
55-64	81.4	2.5	434	68.3	3.8	514	74.3	3.2	948
65+	74.4	3.4	459	56.2	3.9	511	64.8	3.7	970
Total	90.9	2.0	5585	84.7	1.8	5806	87.7	1.9	11391

**Table 13: Percentage of the population using improved drinking water sources, Country, Year**

	Main source of water															Total with safe No. of drinking persons		
	Piped into dwelling	Piped into yard or plot	Public tap	Tube-well/Bore-hole with pump	Pro-ected spring	Pro-ected dug well	Rain-water collec-tion	Bottled water	Unpro-ected spring	Unpro-ected dug well	River or stream	Tanker truck vendor	Cut official pipe	Other	Don't know			Total
Central	79.8	14.0	.7	.0	.1	.0	4.3	.1	.0	.0	.0	.0	.0	.5	.4	100.0	98.9	6778
South Central	44.3	24.8	.4	1.3	1.7	.4	21.9	.0	.3	.0	.0	.0	2.2	2.5	.2	100.0	94.8	3873
West	49.9	31.9	.6	.2	.0	.1	5.7	.3	.0	.2	.8	.9	6.6	1.7	1.1	100.0	88.4	2716
East	32.9	20.9	1.0	1.3	2.8	.2	28.8	.3	2.0	.0	7.5	.0	1.0	1.1	.0	100.0	87.9	2077
South	.0	7.4	2.1	.0	.0	.0	21.3	.0	.0	.5	68.5	.0	.0	.0	.3	100.0	30.8	1518
Urban	74.4	16.2	.5	.5	.6	.2	5.5	.1	.2	.0	.0	.1	.1	1.1	.5	100.0	97.9	9312
Rural	29.2	23.7	1.1	.5	.9	.1	22.3	.1	.5	.2	15.9	.2	3.6	1.3	.4	100.0	77.8	7651
Total	54.0	19.6	.8	.5	.8	.1	13.1	.1	.3	.1	7.2	.1	1.7	1.2	.4	100.0	88.9	1696

3

**Table 14: Percentage of the population using sanitary means of excreta disposal, Country, Year**

	Type of toilet facility									Total	Total with sanitary means of excreta disposal	No. of persons
	Flush to sewage system/ septic tank	Pour flush latrine	Improved pit latrine	Traditional pit latrine	River	Open pit	Other	No facilities/ bush/field	Missing			
Central	87.5	11.5	.2	.0	.0	.0	.5	.0	.4	100.0	99.2	6778
South Central	73.7	25.7	.0	.2	.0	.1	.1	.1	.0	100.0	99.6	3873
West	71.0	24.3	1.5	1.0	.0	.6	.6	.2	.7	100.0	97.8	2716
East	50.9	44.3	.5	1.0	.0	1.7	1.2	.4	.0	100.0	96.7	2077
South	.8	17.4	.0	.6	28.6	1.1	.0	50.8	.6	100.0	18.9	1518
Urban	84.3	14.6	.1	.0	.0	.0	.5	.0	.4	100.0	99.1	9312
Rural	51.5	29.5	.7	.8	5.7	.9	.4	10.3	.2	100.0	82.4	7651
Total	69.5	21.3	.4	.4	2.6	.4	.5	4.7	.3	100.0	91.6	16963

**Table 15: Percentage of under-five children who are severely or moderately undernourished, Country, Year**

	Weight for age		Height for age		Weight for height		Number of children
	Percent below	Percent below	Percent below	Percent below	Percent below	Percent below	
	- 2 SD	-3 SD	- 2 SD	-3 SD	-2 SD	-3 SD	
Male	10.8	2.7	14.2	2.5	3.9	0.8	887
Female	7.1	1.9	11.6	1.3	2.6	0.6	779
Central	5.8	1.4	10.8	1.9	3.3	0.8	609
South Central	8.7	2.5	16.6	1.6	4.6	1.1	350
West	6.1	1.7	11.1	.8	3.9	0.2	228
East	7.4	.5	6.9	.9	1.0	0.2	207
South	20.4	6.0	19.4	3.9	2.9	0.7	272
Urban	6.2	1.3	10.8	1.4	3.3	0.6	816
Rural	11.9	3.3	15.1	2.4	3.3	0.8	849
< 6 months	2.5	.5	3.6	1.0	0.5	0.3	196
6-11 months	8.4	2.9	11.0	3.4	4.3	2.1	141
12-23 months	13.1	4.2	16.8	2.3	5.2	0.7	346
24-35 months	8.6	1.6	12.4	1.6	3.4	0.3	378
36-47 months	9.5	2.5	14.3	1.6	2.9	1.0	315
48-59 months	8.8	1.7	14.8	2.0	2.8	0.7	288
Mother's education							
None	15.5	4.5	15.1	3.7	3.3	0.9	211
Primary	13.2	2.4	16.2	2.2	3.2	0.7	444
Secondary +	5.9	1.8	11.2	1.4	3.4	0.7	1010
<b>Total</b>	<b>9.1</b>	<b>2.3</b>	<b>13.0</b>	<b>1.9</b>	<b>3.3</b>	<b>0.7</b>	<b>1665</b>

**Table 16: Percent of living children by breastfeeding status, Country, Year**

	Percent of children 0-3 months exclusively breastfed		Percent of children 6-9 months receiving breastmilk and solid/semi-solid food		Percent of children 12-15 months breastfed		Percent of children 20-23 months breastfed	
	Percent of children 0-3 months exclusively breastfed	Number of children	Percent of children 6-9 months receiving breastmilk and solid/semi-solid food	Number of children	Percent of children 12-15 months breastfed	Number of children	Percent of children 20-23 months breastfed	Number of children
Male	14.5	77	18.6	59	39.5	66	10.9	48
Female	9.9	69	32.6	49	36.9	72	13.4	52
Urban	13.0	72	23.9	48	31.1	64	18.2	46
Rural	11.7	74	25.8	59	44.2	74	7.1	54
Total	12.4	146	25.0	108	38.1	138	12.2	100

**Table 17: Percentage of households consuming adequately iodized salt, Country, Year**

	Percent of households with no salt	Percent of households in which salt was tested	Percent of households with salt testing		Number of households interviewed
			< 15 PPM	15+ PPM	
Central	0.3	99.6	10.6	89.4	1778
South Central	0.8	98.9	30.9	69.1	938
West	0.7	99.3	26.6	73.4	673
East	1.0	96.8	22.9	77.1	493
South	5.0	89.3	42.9	57.1	403
Urban	0.9	99.5	18.6	81.4	2395
Rural	1.1	96.4	26.3	73.7	1890
<b>Total</b>	<b>1.0</b>	<b>98.1</b>	<b>22.0</b>	<b>78.0</b>	<b>4285</b>

Note: Adequately iodized salt is salt testing 15 PPM (parts per million) or more.

**Table 18: Percent distribution of children aged 6-59 months by whether they have received a high dose Vitamin A supplement in the last 6 months, Country, Year**

	Percent of children who received Vitamin A:			Not sure if received	Never received	Total	Number of children
	Within last 6 months	Prior to last 6 months	Not sure when				
Male	12.2	6.1	0.2	2.2	79.3	100.0	855
Female	15.6	6.6	0.1	1.3	76.4	100.0	757
Central	14.2	8.5	0.3	2.1	74.8	100.0	576
South Central	14.2	5.4	0.0	1.5	78.9	100.0	331
West	17.1	8.9	0.4	1.6	72.0	100.0	246
East	15.1	5.5	0.0	1.5	77.9	100.0	199
South	8.1	0.8	0.0	1.9	89.2	100.0	260
Urban	15.9	8.0	0.3	2.2	73.8	100.0	786
Rural	11.7	4.7	0.1	1.5	82.0	100.0	826
6-11 months	17.1	1.3	0.0	0.6	81.0	100.0	158
12-23 months	21.0	5.2	0.3	1.1	72.4	100.0	366
24-35 months	13.7	9.7	0.0	0.5	76.3	100.0	422
36-47 months	9.6	5.2	0.0	3.2	82.0	100.0	345
48-59 months	8.4	6.9	0.6	3.4	80.7	100.0	321
Mother's education							
None	6.3	3.9	0.5	1.9	87.4	100.0	207
Primary	13.8	7.3	0.5	1.9	76.5	100.0	426
Secondary +	15.3	6.4	0.0	1.7	76.5	100.0	979
<b>Total</b>	<b>13.8</b>	<b>6.3</b>	<b>0.2</b>	<b>1.8</b>	<b>77.9</b>	<b>100.0</b>	<b>1612</b>

**Table 19: Percentage of women with a birth in the last 12 months by whether they received a high dose Vitamin A supplement before the infant was 8 weeks old, Country, Year**

	<b>Received Vitamin A supplement</b>	<b>Not sure if received</b>	<b>Number of women</b>
Central	15.2	0.0	145
South Central	13.3	0.0	75
West	<b>15.6</b>	0.0	45
East	10.5	2.6	38
South	2.4	0.0	84
Urban	12.8	0.0	195
Rural	10.4	0.5	192
Mother's education			
None	8.7	0.0	46
Primary	10.5	0.9	114
Secondary +	12.8	0.0	228
<b>Total</b>	<b>11.6</b>	<b>0.3</b>	<b>387</b>

**Table 20: Percentage of live births in the last 12 months that weighed below 2500 grams at birth, Country, Year**

	Percent of live births		Number of live births
	Below 2500 grams	Weighed at birth	
Central	12.8	86.0	145
South Central	12.7	85.7	75
West	10.2	85.7	45
East	14.4	73.2	38
South	9.8	57.8	84
Urban	12.1	87.4	195
Rural	11.8	69.6	192
Mother's education			
None	11.0	54.2	46
Primary	11.5	70.3	114
Secondary +	12.4	87.6	228
Total	12.0	78.6	387

**Table 21: Percentage of children age 12-23 months immunized against childhood diseases at any time before the survey and before the first birthday, Country, Year**

Percentage of children who received:											
	BCG	DPT1	DPT2	DPT3	Polio 1	Polio2	Polio3	Measles	All	None	No. of children
Vaccinated at any time before the survey											
According to:											
Vaccination card	85.2	80.5	78.6	74.3	84.7	81.8	77.8	56.2	51.7	1.1	314
Mother's report	3.7	5.0	3.0	3.0	3.3	3.3	2.5	4.4	0.8	2.7	52
Either	88.9	85.5	81.6	77.3	88.0	85.1	80.3	60.6	52.5	3.8	366
Vaccinated by 12 months of age	87.1	84.0	79.4	74.5	86.9	83.1	77.2	24.0	19.3	-	366

**Table 22: Percentage of children age 12-23 months currently vaccinated against childhood diseases, Country, Year**

	BCG	DPT 1	DPT 2	DPT 3	Polio 1	Polio 2	Polio 3	Measles	All	None	% with health card	Number of children
Male	88.5	86.1	80.6	76.9	88.1	83.7	79.0	57.7	50.0	2.6	86.9	183
Female	89.1	84.8	82.7	77.8	88.0	86.5	81.6	63.5	54.9	4.9	84.7	183
Central	93.2	88.4	83.0	80.4	87.5	85.7	82.1	63.4	56.2	2.7	83.0	117
South Central	96.4	94.0	89.2	85.7	90.4	88.1	84.7	59.5	53.6	.0	87.9	84
West	95.6	93.5	95.8	93.5	95.8	93.5	93.5	61.2	58.8	.0	95.8	45
East	76.1	75.0	75.0	72.9	75.0	72.9	70.8	52.1	45.8	10.4	75.0	46
South	75.7	72.7	66.2	55.8	89.6	83.1	70.1	62.3	45.5	7.8	88.3	74
Urban	91.2	89.6	85.3	82.8	89.0	87.1	84.7	63.2	56.4	1.8	84.7	171
Rural	86.2	81.9	78.4	72.5	87.3	83.3	76.5	58.3	49.0	5.4	86.8	196
Mother's education												
None	77.6	68.6	61.7	48.4	78.2	73.2	58.3	53.7	35.1	9.9	79.8	58
Primary	86.0	79.9	77.9	72.7	85.2	82.0	76.8	50.5	45.3	5.1	84.0	93
Secondary +	93.5	92.4	88.7	87.2	92.0	89.7	87.8	66.9	60.3	1.5	88.2	214
Total	88.9	85.5	81.6	77.3	88.0	85.1	80.3	60.6	52.5	3.7	85.8	366

**Table 23: Percentage of under-five children with diarrhea in the last two weeks and treatment with ORS or ORT, Country, Year**

	Had diarrhea in last two weeks	Number of children under 5	Children with diarrhea who received:					Number of children with diarrhea
			Breast milk	Gruel	ORS packet	Any recommended treatment	No treatment	
Male	13.5	970	21.2	30.4	34.0	61.3	38.7	131
Female	14.3	860	28.8	23.7	33.2	60.6	39.4	123
Central	12.1	663	23.4	36.4	33.8	61.0	39.0	81
South Central	12.1	370	15.6	26.5	31.3	58.0	42.0	45
West	8.3	270	*	*	*	*	*	22
East	13.2	219	26.7	40.0	16.7	60.0	40.0	29
South	25.2	309	29.6	13.6	45.7	65.4	34.6	78
Urban	11.6	896	21.2	33.3	34.3	61.6	38.4	104
Rural	16.1	934	27.4	22.9	33.1	60.5	39.5	151
< 6 months	14.1	215	58.2	32.4	22.9	74.5	25.5	30
6-11 months	23.4	158	53.6	54.3	37.7	91.7	8.3	37
12-23 months	19.2	366	25.6	33.1	39.6	66.8	33.2	70
24-35 months	12.7	422	9.3	14.8	29.4	46.1	53.9	54
36-47 months	9.8	345	5.7	20.6	23.5	40.7	59.3	34
48-59 months	9.1	321	3.3	3.6	44.9	44.9	55.1	29
Mother's education								
None	22.8	238	28.7	16.0	37.3	56.9	43.1	54
Primary	17.4	496	28.1	25.2	40.0	67.1	32.9	86
Secondary +	10.4	1096	20.6	34.0	27.0	58.3	41.7	114
<b>Total</b>	<b>13.9</b>	<b>1830</b>	<b>24.9</b>	<b>27.2</b>	<b>33.6</b>	<b>61.0</b>	<b>39.0</b>	<b>254</b>

**Table 24: Percentage of under-five children with diarrhea in the last two weeks who took increased fluids and continued to feed during the episode, Country, Year**

	Had diarrhea in last two weeks	Number of children under 5	Children with diarrhea who drank:				Children with diarrhea who ate:				Received increased fluids and continued eating	Number of children with diarrhea
			More	Same/Less	Missing/DK	Total	Somewhat less/same/more	Much less/none	Missing/DK	Total		
Male	13.5	970	31.4	65.5	3.1	100.0	74.0	23.0	3.1	100.0	23.6	131
Female	14.3	860	33.5	62.6	3.9	100.0	67.0	27.5	5.4	100.0	26.0	123
Central	12.1	663	45.5	53.2	1.3	100.0	67.5	31.2	1.3	100.0	33.8	81
South Central	12.1	370	27.2	66.1	6.6	100.0	64.4	29.0	6.6	100.0	17.9	45
West	8.3	270	*	*	*	*	*	*	*	*	*	22
East	13.2	219	43.3	56.7	.0	100.0	60.0	36.7	3.3	100.0	26.7	29
South	25.2	309	23.5	71.6	4.9	100.0	75.3	18.5	6.2	100.0	22.2	78
Urban	11.6	896	43.4	54.5	2.0	100.0	67.7	30.3	2.0	100.0	30.3	104
Rural	16.1	934	24.8	70.7	4.5	100.0	72.6	21.7	5.7	100.0	21.0	151
< 6 months	14.1	215	20.3	66.8	12.9	100.0	74.2	6.6	19.2	100.0	16.9	30
6-11 months	23.4	158	24.6	72.7	2.6	100.0	70.1	27.3	2.6	100.0	21.8	37
12-23 months	19.2	366	26.9	70.3	2.7	100.0	71.8	26.8	1.4	100.0	23.9	70
24-35 months	12.7	422	48.5	51.5	.0	100.0	79.2	20.8	.0	100.0	33.3	54
36-47 months	9.8	345	32.5	64.4	3.1	100.0	58.8	38.1	3.1	100.0	20.9	34
48-59 months	9.1	321	38.3	58.4	3.3	100.0	62.3	31.1	6.6	100.0	27.8	29
Mother's education												
None	22.8	238	28.7	69.4	1.9	100.0	67.6	26.9	5.4	100.0	24.8	54
Primary	17.4	496	28.9	67.7	3.3	100.0	77.3	18.2	4.5	100.0	26.6	86
Secondary +	10.4	1096	36.8	58.9	4.3	100.0	66.9	29.6	3.5	100.0	23.4	114
Total	13.9	1830	32.4	64.1	3.5	100.0	70.6	25.2	4.2	100.0	24.8	254

\* Fewer than 25 cases

**Table 25: Percentage of under-five children with acute respiratory infection in the last two weeks and treatment by health providers, Country, Year**

	Had acute respiratory infection	Number of children under 5	Children with ARI who were taken to							Any appropriate provider	Number of children with ARI
			Family/Friend/Neighbor	Doctor	Specialist	Nurse/Health assistant	Hospital	Other			
Male	4.7	970	4.2	34.9	6.7	14.7	2.1	6.7	54.2	46	
Female	3.7	860	6.3	43.4	3.3	12.0	.0	.0	58.7	32	
Urban	3.7	896	3.1	37.5	9.4	.0	.0	6.2	46.9	33	
Rural	4.7	934	6.5	39.1	2.2	23.9	2.2	2.2	63.0	44	
Total	4.2	1830	5.1	38.4	5.3	13.6	1.2	3.9	56.1	78	

**Table 26: Percentage of children 0-59 months of age reported ill during the last two weeks who received increased fluids and continued feeding, Country, Year**

	Reported illness in last two weeks	Number of children under 5	Children with illness who drank:				Children with illness who ate:				Received increased fluids and continued eating	Number of sick children
			More	Same/Less	Missing/DK	Total	Somewhat less/same/more	Much less/none	Missing/DK	Total		
Male	45.2	970	19.9	78.3	1.8	100.0	75.8	22.2	2.0	100.0	14.9	438
Female	45.7	860	21.9	75.7	2.5	100.0	76.4	20.9	2.7	100.0	16.3	393
Central	40.7	663	25.6	73.3	1.2	100.0	81.4	17.4	1.2	100.0	19.4	270
South Central	47.2	370	17.2	81.1	1.7	100.0	74.3	24.0	1.7	100.0	12.5	174
West	37.4	270	16.5	78.8	4.7	100.0	78.9	16.4	4.7	100.0	11.7	101
East	47.4	219	28.7	70.4	.9	100.0	71.3	26.9	1.9	100.0	20.4	104
South	59.0	309	15.3	81.6	3.2	100.0	71.1	25.3	3.7	100.0	12.1	182
Urban	40.8	896	24.0	74.9	1.1	100.0	79.7	19.1	1.1	100.0	18.0	366
Rural	49.8	934	18.4	78.8	2.9	100.0	73.2	23.5	3.3	100.0	13.6	465
< 6 months	38.6	215	10.0	84.1	6.0	100.0	78.8	12.9	8.3	100.0	8.7	83
6-11 months	50.4	158	15.3	82.3	2.4	100.0	74.1	21.1	4.8	100.0	14.0	79
12-23 months	54.3	366	16.6	81.5	2.0	100.0	74.1	24.4	1.5	100.0	14.0	199
24-35 months	45.8	422	29.5	69.5	1.0	100.0	79.0	20.5	.5	100.0	22.2	194
36-47 months	39.9	345	23.0	76.2	.8	100.0	72.6	26.6	.8	100.0	13.8	138
48-59 months	43.1	321	22.4	74.8	2.8	100.0	77.7	19.5	2.8	100.0	15.2	139
None	55.1	238	20.7	77.7	1.5	100.0	68.8	29.0	2.3	100.0	14.8	131
Mother's education												
Primary	45.2	496	18.2	80.1	1.7	100.0	75.6	21.8	2.6	100.0	15.1	224
Secondary +	43.4	1096	22.1	75.4	2.5	100.0	78.3	19.4	2.3	100.0	16.0	476
Total	45.4	1830	20.8	77.0	2.1	100.0	76.1	21.6	2.3	100.0	15.5	831

**Table 27: Percentage of caretakers of children 0-59 months who know at least 2 signs for seeking care immediately, Country, Year**

	Knows child should be taken to health facility if child:							Number of caretakers	
	Not able to drink /breastfeed	Becomes sicker	Develops a fever	Has fast breathing	Has difficult breathing	Has blood in stool	Is drinking poorly		Knows at least two signs
Central	.0	8.7	68.0	.9	6.9	1.1	.6	10.3	663
South Central	.3	15.2	68.7	2.6	8.1	4.5	3.7	17.6	370
West	10.4	16.1	74.0	13.3	17.2	11.4	11.4	21.9	270
East	.9	10.1	64.5	2.6	5.3	.9	.4	9.6	219
South	14.0	22.4	91.6	21.1	24.2	14.6	14.0	34.2	309
Urban	.4	10.2	69.0	1.4	6.3	1.5	1.2	11.2	896
Rural	7.6	16.8	76.1	11.8	16.3	9.4	8.8	23.4	934
Mother's education									
None	6.9	19.1	83.1	11.3	17.4	7.2	7.7	26.8	238
Primary	7.7	17.1	74.0	11.4	16.0	9.7	8.3	20.1	496
Secondary +	1.8	10.8	69.7	3.6	8.0	3.3	3.1	14.2	1096
Total	4.0	13.6	72.6	6.7	11.4	5.6	5.1	17.4	1830

**Table 28: Percentage of children 0-59 months of age who slept under an insecticide-impregnated bednet during the previous night, Country, Year**

	Slept under a bednet			Number of children	Bednet treated		Children who slept under a bednet
	Yes	No	DK/missing		Yes	No	
Male	70.7	23.4	5.9	180	3.8	96.2	128
Female	72.6	20.2	7.1	161	5.7	94.3	117
< 6 months	90.7	7.4	1.9	52	6.1	93.9	47
6-11 months	81.8	15.2	3.0	32	3.7	96.3	26
12-23 months	68.6	23.3	8.1	82	8.5	91.5	57
24-35 months	71.0	23.2	5.8	66	4.1	95.9	47
36-47 months	63.6	25.5	10.9	53	2.9	97.1	34
48-59 months	61.0	32.2	6.8	57	.0	100.0	35
Total	71.6	21.9	6.5	341	4.7	95.3	244

**Table 29: Percentage of children 0-59 months of age who were ill with fever in the last two weeks who received anti-malarial drugs, Country, Year**

	Had a fever in last two weeks	Number of children under 5	Children with a fever who were treated with:					Don't know	Any appropriate anti-malarial drug*	Number of children
			Para-etamol	Chloro-quine	Fansidar	Other				
Male	36.4	970	57.2	54.1	0.3	28.6	1.4	54.4	353	
Female	33.7	860	55.0	52.4	0.3	21.4	1.6	52.8	290	
Central	33.5	663	59.0	55.4	0.0	25.7	0.9	55.4	222	
South Central	34.9	370	50.4	44.2	0.0	32.6	2.3	44.2	129	
West	36.7	270	55.6	53.5	0.0	21.2	1.0	53.5	99	
East	34.2	219	56.4	52.8	1.3	29.3	2.7	54.7	75	
South	38.2	309	57.6	59.3	0.8	18.1	1.7	60.2	118	
Urban	33.3	896	60.3	61.4	0.3	21.8	1.0	61.6	299	
Rural	36.8	934	52.7	46.5	0.3	28.5	2.0	46.8	344	
<6 months	32.2	215	56.3	49.1	0.1	21.9	1.0	49.1	69	
6-11 months	50.6	158	60.4	59.7	0.1	26.5	1.3	59.7	80	
12-23 months	50.8	366	54.9	53.7	0.4	28.5	1.9	54.4	186	
24-35 months	32.1	422	55.5	50.9	0.7	28.1	1.3	51.6	136	
36-47 months	31.0	345	58.9	48.8	0.1	21.7	1.9	48.8	107	
48-59 months	20.2	321	50.9	61.6	0.0	18.5	1.6	61.6	65	
Mother's education										
None	37.1	238	76.0	52.2	0.1	21.9	4.5	52.2	88	
Primary	34.9	496	74.5	56.0	0.6	26.4	2.3	56.6	173	
Secondary +	25.0	1096	60.2	73.0	0.4	35.8	0.7	73.4	274	
Total	35.1	1830	56.1	53.3	0.3	25.4	1.5	53.7	643	

**Table 30: Percentage of women aged 15-49 who know the main ways of preventing HIV transmission, Country, Year**

	Percent who know transmission can be prevented by:							Number of women
	Heard of AIDS	Have only one faithful uninfected sex partner	Using a condom every time	Abstaining from sex	Knows all three ways	Knows at least one way	Doesn't know any way	
Central	97.0	76.3	76.9	53.7	46.8	83.8	16.2	1587
South Central	93.1	63.3	63.5	45.4	37.3	72.1	27.9	953
West	92.7	53.1	46.0	37.5	27.3	60.8	39.2	627
East	91.1	61.6	59.4	43.3	36.3	66.7	33.3	437
South	82.7	25.0	22.2	20.9	14.9	29.4	70.6	360
Urban	96.5	73.9	73.4	52.1	45.0	81.5	18.5	2196
Rural	89.6	50.0	47.6	36.2	28.0	56.9	43.1	1769
15-19	92.0	58.5	57.1	42.9	33.8	66.4	33.6	818
20-24	94.3	60.8	63.1	42.5	33.4	71.6	28.4	631
25-29	94.2	66.5	63.9	46.1	39.4	72.6	27.4	593
30-34	95.1	69.0	64.4	47.4	41.1	73.6	26.4	599
35-39	94.4	66.9	66.1	48.8	42.2	74.3	25.7	531
40-44	93.6	65.8	63.8	47.3	39.4	72.8	27.2	441
45-49	89.2	54.6	54.5	39.8	33.2	60.6	39.4	351
Education								
None	80.7	17.3	14.9	10.8	6.4	21.6	78.4	266
Primary	87.7	40.4	38.1	29.8	22.0	47.4	52.6	1050
Secondary +	97.0	76.9	76.1	54.5	46.6	84.6	15.4	2649
<b>Total</b>	<b>93.4</b>	<b>63.2</b>	<b>61.9</b>	<b>45.0</b>	<b>37.4</b>	<b>70.5</b>	<b>29.5</b>	<b>3965</b>

**Table 31: Percentage of women aged 15-49 who correctly identify misconceptions about HIV/AIDS, Country, Year**

	Percent who know that:							Number of women
	AIDS cannot be transmitted by:			A healthy looking person can be infected	Knows all three misconceptions	Knows at least one misconception	Doesn't correctly identify any misconception	
	Heard of AIDS	Supernatural means	Mosquito bites					
Central	97.0	71.9	63.2	82.7	52.0	90.8	9.2	1587
South Central	93.1	59.7	48.9	71.7	36.2	82.4	17.6	953
West	92.7	49.3	38.8	68.0	28.9	78.2	21.8	627
East	91.1	52.9	45.6	63.3	29.3	77.5	22.5	437
South	82.7	26.3	18.0	46.1	13.4	53.9	46.1	360
Urban	96.5	68.8	59.6	80.9	48.3	89.2	10.8	2196
Rural	89.6	47.1	37.8	61.5	26.4	73.0	27.0	1769
15-19	92.0	55.9	46.3	68.8	32.2	81.2	18.8	818
20-24	94.3	60.0	49.4	74.4	39.1	83.6	16.4	631
25-29	94.2	62.6	55.6	74.8	44.0	83.2	16.8	593
30-34	95.1	62.7	52.6	73.6	41.2	84.0	16.0	599
35-39	94.4	63.6	54.4	72.7	43.7	82.5	17.5	531
40-44	93.6	55.8	47.8	73.7	37.4	81.9	18.1	441
45-49	89.2	51.0	40.7	67.3	31.8	74.7	25.3	351
Education								
None	80.7	17.1	11.2	37.3	7.7	45.1	54.9	266
Primary	87.7	35.0	26.7	53.9	15.8	65.3	34.7	1050
Secondary +	97.0	73.0	63.0	83.0	50.6	92.3	7.7	2649
<b>Total</b>	<b>93.4</b>	<b>59.2</b>	<b>49.9</b>	<b>72.2</b>	<b>38.5</b>	<b>82.0</b>	<b>18.0</b>	<b>3965</b>

**Table 32: Percentage of women aged 15-49 who correctly identify means of HIV transmission from mother to child, Country, Year**

	Know AIDS can be transmitted from mother to child	Percent who know AIDS can be transmitted:				Did not know any specific way	Number of women
		During pregnancy	At delivery	Through breastmilk	Knows all three		
Central	86.1	80.7	57.3	46.2	32.8	15.6	1587
South Central	75.4	70.5	50.4	45.3	31.6	26.3	953
West	70.0	64.2	46.4	45.6	31.7	32.8	627
East	73.5	70.1	50.5	48.0	34.8	27.2	437
South	42.3	38.9	33.2	35.8	28.6	59.0	360
Urban	84.0	78.3	55.6	46.6	32.6	17.9	2196
Rural	65.2	61.2	45.2	43.3	31.7	36.3	1769
15-19	70.9	65.1	44.0	45.2	30.5	31.6	818
20-24	76.5	71.0	49.5	46.0	31.2	24.4	631
25-29	78.2	73.1	54.8	41.0	30.2	23.5	593
30-34	78.9	74.3	54.5	46.3	33.9	22.8	599
35-39	78.1	72.4	55.7	45.0	34.2	24.0	531
40-44	77.8	73.7	52.8	50.3	35.7	24.0	441
45-49	68.5	66.4	47.7	42.4	30.9	32.7	351
Education							
None	43.8	40.6	33.8	36.8	29.9	57.3	266
Primary	58.2	54.0	40.9	42.0	30.7	43.9	1050
Secondary +	85.7	80.3	56.6	47.2	33.0	15.9	2649
<b>Total</b>	<b>75.6</b>	<b>70.7</b>	<b>51.0</b>	<b>45.1</b>	<b>32.2</b>	<b>26.1</b>	<b>3965</b>

**Table 33: Percentage of women aged 15-49 who express a discriminatory attitude towards people with HIV/AIDS, Country, Year**

	Percent of women who:				Number of women
	Believe that a teacher with HIV should not be allowed to work	Would not buy food from a person with HIV/AIDS	Agree with at least one discriminatory statement	Agree with neither discriminatory statement	
Central	63.8	15.2	65.0	35.0	1587
South Central	47.5	11.1	48.9	51.1	953
West	40.0	12.7	41.8	58.2	627
East	40.8	11.9	42.0	58.0	437
South	25.3	21.6	29.6	70.4	360
Urban	60.3	14.2	61.6	38.4	2196
Rural	37.4	13.9	39.5	60.5	1769
15-19	49.0	18.2	51.6	48.4	818
20-24	51.4	14.6	53.0	47.0	631
25-29	51.2	13.2	53.3	46.7	593
30-34	51.2	12.8	51.8	48.2	599
35-39	51.5	12.0	52.4	47.6	531
40-44	50.6	13.6	51.6	48.4	441
45-49	43.9	10.4	45.8	54.2	351
Education					
None	21.9	14.8	25.8	74.2	266
Primary	25.7	9.0	27.3	72.7	1050
Secondary +	62.6	16.0	64.0	36.0	2649
<b>Total</b>	<b>50.1</b>	<b>14.0</b>	<b>51.7</b>	<b>48.3</b>	<b>3965</b>

**Table 34: Percentage of women aged 15-49 who have sufficient knowledge of HIV/AIDS transmission, Country, Year**

	Heard of AIDS	Know 3 ways to prevent HIV transmission	Correctly identify 3 misconceptions about HIV transmission	Have sufficient knowledge	Number of women
Central	97.0	46.8	52.0	30.3	1587
South	93.1	37.3	36.2	21.2	953
Central					
West	92.7	27.3	28.9	14.9	627
East	91.1	36.3	29.3	15.9	437
South	82.7	14.9	13.4	7.5	360
Urban	96.5	45.0	48.3	28.4	2196
Rural	89.6	28.0	26.4	14.1	1769
15-19	92.0	33.8	32.2	16.6	818
20-24	94.3	33.4	39.1	19.5	631
25-29	94.2	39.4	44.0	26.4	593
30-34	95.1	41.1	41.2	24.2	599
35-39	94.4	42.2	43.7	27.3	531
40-44	93.6	39.4	37.4	22.7	441
45-49	89.2	33.2	31.8	18.9	351
Education					
None	80.7	6.4	7.7	2.1	266
Primary	87.7	22.0	15.8	9.3	1050
Secondary +	97.0	46.6	50.6	29.1	2649
<b>Total</b>	<b>93.4</b>	<b>37.4</b>	<b>38.5</b>	<b>22.0</b>	<b>3965</b>

**Table 35: Percentage of women aged 15-49 who know where to get an AIDS test and who have been tested, Country, Year**

	Know a place to get tested	Have been tested	If tested, have been told result	Number of women
Central	72.6	14.4	87.9	1587
South Central	61.5	10.3	88.7	953
West	46.3	9.2	75.9	627
East	55.0	9.6	84.4	437
South	32.0	7.7	66.7	360
Urban	69.3	13.4	88.4	2196
Rural	48.8	9.1	78.6	1769
15-19	55.3	3.1	64.9	818
20-24	65.9	11.6	88.1	631
25-29	63.8	14.3	88.5	593
30-34	60.1	16.6	89.0	599
35-39	59.3	13.6	79.4	531
40-44	61.5	12.7	76.5	441
45-49	54.2	12.5	95.1	351
Education				
None	19.3	7.5	76.8	266
Primary	42.8	10.1	80.9	1050
Secondary +	71.1	12.4	86.8	2649
<b>Total</b>	<b>60.1</b>	<b>11.5</b>	<b>84.9</b>	<b>3965</b>

**Table 36: Percentage of married or in union women aged 15-49 who are using (or whose partner is using) a contraceptive method, Country, Year**

	Percent of married or in–union women who are using:												Total	Any modern method	Any traditional method	Any method	Number of currently married women
	No method	Female sterilization	Male sterilization	Pill	IUD	Injections	Condom	Dia-phragm/foam/jelly	LAM	Periodic abstinence	Withdrawal	Other					
Central	47.0	10.9	.2	30.2	2.5	3.4	4.5	.0	.2	.5	.2	.3	100.0	51.7	1.2	53.0	946
South Central	56.1	13.5	.0	23.2	1.7	2.0	1.6	.0	.0	.9	.7	.2	100.0	42.0	1.8	43.9	625
West	48.9	8.3	.0	33.3	1.3	4.5	1.9	.0	.0	.4	.7	.7	100.0	49.4	1.8	51.1	461
East	61.3	9.0	.0	22.7	2.0	1.3	1.7	.3	.0	.7	.3	.7	100.0	37.0	1.7	38.7	279
South	94.5	.0	.0	4.0	.0	.4	.0	.0	.0	.4	.4	.4	100.0	4.3	1.2	5.5	235
Urban	48.5	10.9	.2	29.8	2.3	3.3	3.7	.0	.2	.5	.3	.5	100.0	50.1	1.4	51.5	1369
Rural	63.7	8.7	.0	21.2	1.3	2.1	1.3	.1	.0	.7	.6	.3	100.0	34.6	1.7	36.3	1176
< 20 years	74.4	.5	.0	19.0	.6	.6	2.3	.0	.0	.5	1.0	1.0	100.0	23.0	2.6	25.6	177
20-24 years	57.1	.2	.0	36.4	.8	1.8	2.2	.0	.3	.5	.5	.3	100.0	41.4	1.5	42.9	390
25-49 years	53.5	12.6	.1	24.4	2.1	3.1	2.7	.0	.1	.6	.4	.4	100.0	45.0	1.4	46.5	1979
Education																	
None	88.6	4.5	.0	5.9	.5	.5	.0	.0	.0	.0	.0	.0	100.0	11.4	.0	11.4	198
Primary	65.1	10.4	.0	17.0	.9	3.8	1.3	.0	.0	.3	.7	.5	100.0	33.5	1.5	34.9	740
Secondary +	47.1	10.3	.1	32.3	2.4	2.6	3.5	.1	.1	.8	.4	.4	100.0	51.2	1.7	52.9	1607
Total	55.5	9.9	.1	25.8	1.8	2.7	2.6	.0	.1	.6	.5	.4	100.0	42.9	1.5	44.5	2545

**Table 37: Percentage of mothers with a birth in the last 12 months protected against neonatal tetanus, Country, Year**

	Percent of mothers with a birth in the last 12 months who:			Protected against tetanus	Number of mothers
	Received at least 2 doses, last within 3 years	Received at least 3 doses, last within 10 years	Received at least 5 doses during lifetime		
Central	72.4	5.5	0.0	77.9	145
South Central	72.0	8.0	0.0	80.0	75
West	86.7	2.2	0.0	88.9	45
East	68.4	5.3	0.0	73.7	38
South	61.9	4.8	1.2	67.9	84
Urban	76.4	1.5	0.0	77.9	195
Rural	66.1	9.4	0.5	76.0	192
Education					
None	69.6	8.7	2.2	80.4	46
Primary	78.1	10.5	0.0	88.6	114
Secondary +	68.0	2.2	0.0	70.2	227
<b>Total</b>	<b>71.3</b>	<b>5.4</b>	<b>0.3</b>	<b>77.0</b>	<b>387</b>

**Table 38: Percent distribution of women aged 15-49 with a birth in the last year by type of personnel delivering antenatal care, Country, Year**

	Person delivering antenatal care						Total	Any skilled personnel	Number of women
	Doctor	Nurse	Midwife	Health assistant	Traditional birth attendant	No antenatal care received			
Central	75.7	2.9	9.6	.0	11.0	.7	100.0	88.2	145
South							100.0	95.1	75
Central	71.8	5.7	17.7	.0	4.9	.0	100.0	80.9	45
West	56.9	9.1	15.0	.0	19.1	.0	100.0	87.8	38
East	56.1	9.8	22.0	.0	12.2	.0	100.0	87.8	38
South	7.8	8.9	5.6	73.3	4.4	.0	100.0	22.2	84
Urban	71.0	6.0	13.1	.0	9.3	.5	100.0	90.2	195
Rural	41.1	6.3	11.1	31.9	9.7	.0	100.0	58.5	192
Education									
None	15.2	10.5	10.8	57.1	6.4	.0	100.0	36.5	46
Primary	42.5	10.4	9.4	28.4	9.3	.0	100.0	62.3	114
Secondary +	71.2	3.2	13.7	1.2	10.2	.5	100.0	88.1	227
Total	56.2	6.1	12.1	15.8	9.5	.3	100.0	74.4	387

**Table 39: Percent distribution of women aged 15-49 with a birth in the last year by type of personnel assisting at delivery, Country, Year**

	Person assisting at delivery						Total	Any skilled personnel	Number of women
	Doctor	Nurse	Midwife	Health assistant	Traditional birth attendant	Other/missing			
Central	33.1	16.9	42.6	.0	.0	7.4	100.0	92.6	145
South Central	29.8	20.4	48.5	.0	.0	1.2	100.0	98.8	75
West	32.3	9.1	41.6	.0	2.0	15.0	100.0	83.0	45
East	29.3	9.8	48.8	.0	2.4	9.8	100.0	87.8	38
South	7.8	3.3	10.0	46.7	26.7	5.6	100.0	21.1	84
Urban	31.7	16.4	45.4	.0	.0	6.6	100.0	93.4	195
Rural	21.3	9.7	29.0	20.3	12.6	7.2	100.0	59.9	192
Education									
None	15.2	2.3	16.9	28.5	28.5	8.5	100.0	34.4	46
Primary	19.8	17.5	28.5	20.3	8.1	5.8	100.0	65.7	114
Secondary +	32.2	13.0	45.7	1.2	.8	7.1	100.0	90.8	228
Total	26.5	13.1	37.2	10.1	6.2	6.9	100.0	76.8	387

**Table 40: Percent distribution of children aged 0-59 months by whether birth is registered and reasons for non-registration, Country, Year**

	Birth is registered	DK if birth registered	Birth is not registered because:						Total	No. of children
			Costs too much	Must travel too far	Didn't know it should be registered	Doesn't know where to register	Other	Reason DK or Missing		
Male	93.8	4.3	.1	.3	.1	.1	.7	.5	100.0	970
Female	94.7	4.0	.0	.1	.2	.1	.3	.5	100.0	860
Central	95.1	2.7	.2	.2	.3	.2	.6	.8	100.0	663
South Central	94.9	3.2	.0	.6	.0	.3	1.1	.0	100.0	370
West	85.6	13.3	.0	.0	.4	.0	.0	.7	100.0	270
East	96.5	2.2	.0	.4	.0	.0	.9	.0	100.0	219
South	97.5	1.9	.0	.0	.0	.0	.0	.6	100.0	309
Urban	93.5	4.2	.1	.4	.2	.2	.7	.7	100.0	896
Rural	95.0	4.1	.0	.1	.1	.0	.4	.3	100.0	934
< 6 months	90.7	4.6	.0	.9	.0	.0	2.3	1.4	100.0	215
6-11 months	96.7	2.0	.0	.0	.0	.0	.6	.7	100.0	158
12-23 months	95.6	2.7	.0	.3	.3	.0	.8	.3	100.0	366
24-35 months	93.3	5.4	.2	.0	.2	.0	.2	.5	100.0	422
36-47 months	93.6	4.7	.0	.3	.3	.6	.0	.6	100.0	345
48-59 months	95.7	4.3	.0	.0	.0	.0	.0	.0	100.0	321
Mother's education										
None	92.0	2.9	.0	.8	.9	.9	1.3	1.2	100.0	238
Primary	92.6	5.6	.2	.4	.2	.0	.8	.2	100.0	496
Secondary +	95.5	3.8	.0	.0	.0	.0	.3	.5	100.0	1096
<b>Total</b>	<b>94.2</b>	<b>4.2</b>	<b>.1</b>	<b>.2</b>	<b>.2</b>	<b>.1</b>	<b>.6</b>	<b>.5</b>	<b>100.0</b>	<b>1830</b>

**Table 41: Percentage of children 0-14 years of age in households not living with a biological parent, Country, Year**

	Living with both parents	Living with neither parent				Living with mother only		Living with father only		Impossible to determine	Total	Not living with a biological parent	One or both parents dead	Number of children
		Father M only live	Mother only alive	Both are alive	Both are dead	Father alive	Father dead	Mother alive	Mother dead					
Male	64.5	.6	.2	5.9	.2	20.5	2.3	2.2	.3	3.4	100.0	6.9	3.5	2785
Female	63.0	.7	.4	6.4	.2	22.2	2.1	1.4	.3	3.3	100.0	7.7	3.7	2601
Central	61.5	.7	.4	5.4	.1	24.9	2.6	2.2	.4	1.9	100.0	6.6	4.1	1954
South	75.0	.4	.2	4.1	.2	14.2	2.7	1.8	.1	1.4	100.0	4.9	3.6	1185
Central														
West	71.4	.6	.1	5.7	.6	9.3	1.2	1.9	.5	8.8	100.0	7.0	3.0	816
East	61.7	.7	.1	8.1	.1	22.4	2.3	2.2	.3	2.2	100.0	9.1	3.5	705
South	45.0	.7	.9	10.1	.1	36.1	1.3	.4	.0	5.4	100.0	11.8	3.0	726
Urban	64.2	.7	.3	5.4	.2	21.4	2.7	2.1	.4	2.7	100.0	6.5	4.2	2698
Rural	63.3	.6	.4	6.9	.2	21.2	1.7	1.5	.2	4.0	100.0	8.1	3.1	2688
0-4 years	68.5	.2	.2	3.7	.1	22.5	.9	1.1	.1	2.7	100.0	4.2	1.5	1937
5-9 years	63.8	.6	.2	7.1	.1	22.1	2.4	1.4	.1	2.4	100.0	8.0	3.3	1673
10-14 years	58.6	1.1	.6	7.9	.4	19.3	3.4	2.9	.7	4.9	100.0	10.1	6.2	1776
Total	63.8	.6	.3	6.2	.2	21.3	2.2	1.8	.3	3.3	100.0	7.3	3.6	5386

**Table 42: Percentage of children 5-14 years of age who are currently working, Country, Year**

	Paid work	Unpaid work	Domestic work		Currently working	No. of children
			< 4 hours/day	4 or more hours/day		
Male	2.0	4.7	49.4	25.8	31.7	1765
Female	1.3	4.1	56.4	25.0	29.7	1685
5-9 years	0.5	1.7	46.9	18.3	20.3	1673
10-14 years	2.8	7.0	58.4	32.1	40.4	1776
Central	1.8	3.1	53.9	19.3	23.8	1243
South Central	1.0	5.6	51.1	30.4	36.1	802
West	1.7	3.7	52.9	18.8	24.0	533
East	1.5	5.6	50.3	31.0	37.3	467
South	2.7	5.6	55.7	36.4	42.5	404
Urban	1.8	3.3	54.1	20.1	24.6	1740
Rural	1.5	5.5	51.6	30.8	36.9	1709
Total	1.7	4.4	52.8	25.4	30.7	3449