


**WHO/UNICEF
Review of National Immunization Coverage
1980-2008**

Chile

July, 2009

Chile

BCG (1980-2008)



Description of trend

Trend follows WHO officially reported data. No survey data available.

Data presented in chart

Year	WHO/ UNICEF estimate (%)	Reported to:*		Government official estimate (%)	Reported doses administered (%)**	Survey data (%)***	
		WHO (%)	UNICEF (%)			Survey 12-23 months	Survey <12 months
1980	96	96	96	96	96		
1981	96	96	96	96	96		
1982	94	94	94	94	94		
1983	92	92	92	92	92		
1984	95	95	95	95	95		
1985	96	96	96	96	96		
1986	96	96	96	96	96		
1987	97	97	97	97	97		
1988	98	98	98	98	98		
1989	97	97	97	97	97		
1990	94	94	94	94	94		
1991	99	100	100	100	100		
1992	99	100	100	100	100		
1993	97	97	97	97	97		
1994	99	99	99	99	99		
1995	97	97	97	97	97		
1996	99	99	99	99	99		
1997	92	92	92	92	92		
1998	92	92	92	92	92		
1999	95	95	95	95	95		
2000	93	93	93	93	93		
2001	98	98	98	98	98		
2002	96	96	96	96	96		
2003	95	95	95	95	95		
2004	95	95	95	95	95		
2005	95	95	95	95	95		
2006	98	98	98	98	98		
2007	98	98	98	98	98		
2008	98						


*Prior to 1998 national reports to WHO/UNICEF did not specify whether information was derived from administrative records, surveys or other sources.

**Coverage based on registration of doses administered by health care providers.

***In case more than one survey was implemented in a certain year the highest value is presented. Details of all data are presented in the second section of this report.

Chile

DTP1 (1980-2008)



Description of trend

WHO and UNICEF began requesting data on DTP1 coverage in 2001 and have received national reports reflecting DTP1 coverage from 2001 onward. The DTP1 estimates from 2000-2004 are based on these reports. For years prior to 2000 the estimates are derived from the WHO/UNICEF estimates of DTP3 and the relationship between the levels of DTP3 coverage and the drop-out between DTP1 and DTP3. This relationship results from an analysis of 282 surveys conducted in 101 countries which were published between 1980 and 2004.

Data presented in chart

Year	WHO/ UNICEF estimate (%)	Reported to:*		Government official estimate (%)	Reported doses administered (%)**	Survey data (%)***	
		WHO (%)	UNICEF (%)			Survey 12-23 months	Survey <12 months
1980	97						
1981	99						
1982	99						
1983	96						
1984	99						
1985	99						
1986	96						
1987	97						
1988	98						
1989	98						
1990	98						
1991	97						
1992	96						
1993	96						
1994	97						
1995	97						
1996	97						
1997	96						
1998	96						
1999	98						
2000	98	98	98	98	98		
2001	98	100	100		100		
2002	98						
2003	98	93	93		95		
2004	94	94	94	94	94		
2005	92	92	92	92	92		
2006	95	95	95	95	95		
2007	98	98	98	98	98		
2008	98						


*Prior to 1998 national reports to WHO/UNICEF did not specify whether information was derived from administrative records, surveys or other sources.

**Coverage based on registration of doses administered by health care providers.

***In case more than one survey was implemented in a certain year the highest value is presented. Details of all data are presented in the second section of this report.

Chile

DTP3 (1980-2008)



Description of trend

Trend follows WHO officially reported data. No survey data available.

Data presented in chart

Year	WHO/ UNICEF estimate (%) —	Reported to:*		Government official estimate (%) ○	Reported doses administered (%)** ✕	Survey data (%)***	
		WHO (%) □	UNICEF (%) ■			Survey 12-23 months 	Survey <12 months +
1980	93	93	93	93	93		
1981	99	100	100	100	100		
1982	99	100	100	100	100		
1983	89	89	89	89	89		
1984	99	100	100	100	100		
1985	99	99	99	99	99		
1986	90	90	90	90	90		
1987	93	93	93	93	93		
1988	96	96	96	96	96		
1989	95	95	95	95	95		
1990	95	95	95	95	95		
1991	94	94	94	94	94		
1992	90	90	90	90	90		
1993	90	90	90	90	90		
1994	94	94	94	94	94		
1995	94	94	94	94	94		
1996	91	91	91	91	91		
1997	90	90	90	90	90		
1998	90	90	90	90	90		
1999	95	95	95	95	95		
2000	91	91	91	91	91		
2001	97	97	97	97	97		
2002	97	97	97	97	97		
2003	96	96	96	96	96		
2004	93	93	93	93	93		
2005	91	91	91	91	91		
2006	94	94	94	94	94		
2007	96	96	96	96	96		
2008	96						


*Prior to 1998 national reports to WHO/UNICEF did not specify whether information was derived from administrative records, surveys or other sources.

**Coverage based on registration of doses administered by health care providers.

***In case more than one survey was implemented in a certain year the highest value is presented. Details of all data are presented in the second section of this report.

Chile

Pol3 (1980-2008)



Description of trend

Trend follows WHO officially reported data. No survey data available.

Data presented in chart

Year	WHO/ UNICEF estimate (%) —	Reported to:*		Government official estimate (%) ○	Reported doses administered (%)** ✕	Survey data (%)***	
		WHO (%) □	UNICEF (%) ■			Survey 12-23 months 	Survey <12 months +
1980	93	93	93	93	93		
1981	99	100	100	100	100		
1982	99	100	100	100	100		
1983	89	89	89	89	89		
1984	99	100	100	100	100		
1985	99	99	99	99	99		
1986	90	90	90	90	90		
1987	93	93	93	93	93		
1988	96	96	96	96	96		
1989	95	95	95	95	95		
1990	95	95	95	95	95		
1991	94	94	94	94	94		
1992	90	90	90	90	90		
1993	90	90	90	90	90		
1994	94	94	94	94	94		
1995	94	94	94	94	94		
1996	91	91	91	91	91		
1997	90	90	90	90	90		
1998	90	90	90	90	90		
1999	95	95	95	95	95		
2000	91	91	91	91	91		
2001	96	96	96	96	96		
2002	97	97	97	97	97		
2003	96	96	96	96	96		
2004	94	94	94	94	94		
2005	92	92	92	92	92		
2006	94	94	94	94	94		
2007	95	95	95	95	95		
2008	95						


*Prior to 1998 national reports to WHO/UNICEF did not specify whether information was derived from administrative records, surveys or other sources.

**Coverage based on registration of doses administered by health care providers.

***In case more than one survey was implemented in a certain year the highest value is presented. Details of all data are presented in the second section of this report.

Chile

MCV (1980-2008)



Description of trend

Trend follows WHO officially reported data . No survey data available.

Data presented in chart

Year	WHO/ UNICEF estimate (%) —	Reported to:*		Government official estimate (%) ○	Reported doses administered (%)** ✕	Survey data (%)***	
		WHO (%) □	UNICEF (%) ■			Survey 12-23 months 	Survey <12 months +
1980	94	94	94	94	94		
1981	95	95	95	95	95		
1982	96	96	96	96	96		
1983	95	95	95	95	95		
1984	94	94	94	94	94		
1985	92	92	92	92	92		
1986	91	91	91	91	91		
1987	92	92	92	92	92		
1988	95	95	95	95	95		
1989	92	92	92	92	92		
1990	97	97	97	97	97		
1991	99	99	99	99	99		
1992	89	89	89	89	89		
1993	94	94	94	94	94		
1994	96	96	96	96	96		
1995	97	97	97	97	97		
1996	93	93	93	93	93		
1997	96	96	96	96	96		
1998	98	98	98	98	98		
1999	94	94	94	94	94		
2000	97	97	97	97	97		
2001	96	96	96	96	96		
2002	95	95	95	95	95		
2003	96	96	96	96	96		
2004	94	94	94	94	94		
2005	90	90	90	90	90		
2006	91	91	91	91	91		
2007	92	92	92	92	92		
2008	92						


*Prior to 1998 national reports to WHO/UNICEF did not specify whether information was derived from administrative records, surveys or other sources.

**Coverage based on registration of doses administered by health care providers.

***In case more than one survey was implemented in a certain year the highest value is presented. Details of all data are presented in the second section of this report.

Chile

HepB3 (1980-2008)



Description of trend

Hepatitis B introduced in 2005. No survey data available. Estimate based on reported data.

Data presented in chart

Year	WHO/ UNICEF estimate (%)	Reported to:*		Government official estimate (%)	Reported doses administered (%)**	Survey data (%)***	
		WHO (%)	UNICEF (%)			Survey 12-23 months	Survey <12 months
1980							
1981							
1982							
1983							
1984							
1985							
1986							
1987							
1988							
1989							
1990							
1991							
1992							
1993							
1994							
1995							
1996							
1997							
1998							
1999							
2000							
2001							
2002							
2003							
2004							
2005							
2006	94	94	94	94	94		
2007	96	96	96	96	96		
2008	96						


*Prior to 1998 national reports to WHO/UNICEF did not specify whether information was derived from administrative records, surveys or other sources.

**Coverage based on registration of doses administered by health care providers.

***In case more than one survey was implemented in a certain year the highest value is presented. Details of all data are presented in the second section of this report.

Chile

Hib3 (1980-2008)



Description of trend

Hib vaccine introduced in 1996. Trend follows WHO officially reported data. No survey data available.

Data presented in chart

Year	WHO/ UNICEF estimate (%)	Reported to:*		Government official estimate (%)	Reported doses administered (%)**	Survey data (%)***	
		WHO (%)	UNICEF (%)			Survey 12-23 months	Survey <12 months
1980							
1981							
1982							
1983							
1984							
1985							
1986							
1987							
1988							
1989							
1990							
1991							
1992							
1993							
1994							
1995							
1996	91	91	91	91	91		
1997	90	90	90	90	90		
1998	90	90	90	90	90		
1999	95	95	95	95	95		
2000	91	91	91	91	91		
2001	97	97	97	97	97		
2002	97	97	97	97	97		
2003	96	96	96	96	96		
2004	93	93	93	93	93		
2005	91	91	91	91	91		
2006	94	94	94	94	94		
2007	96	96	96	96	96		
2008	96						

*Prior to 1998 national reports to WHO/UNICEF did not specify whether information was derived from administrative records, surveys or other sources.

**Coverage based on registration of doses administered by health care providers.

***In case more than one survey was implemented in a certain year the highest value is presented. Details of all data are presented in the second section of this report.